

References

- [1] J. M. Link et al. A measurement of lifetime differences in the neutral D meson system. *Phys. Lett.*, B485:62–70, 2000.
- [2] J. M. Link et al. Measurements of the Σ_c^0 and Σ_c^{++} mass splittings. *Phys. Lett.*, B488:218–224, 2000.
- [3] J. M. Link et al. Search for CP violation in D^0 and D^+ decays. *Phys. Lett.*, B491:232–239, 2000.
- [4] J. M. Link et al. A study of the decay $D^0 \rightarrow K^+ \pi^-$. *Phys. Rev. Lett.*, 86:2955–2958, 2001.
- [5] J. M. Link et al. A measurement of branching ratios of D^+ and D_s^+ hadronic decays to four-body final states containing a K_S^0 . *Phys. Rev. Lett.*, 87:162001, 2001.
- [6] J. M. Link et al. Measurement of the relative branching ratio $\text{BR}(\Xi_c^+ \rightarrow p K^- \pi^+)/\text{BR}(\Xi_c^+ \rightarrow \Xi^- \pi^+ \pi^+)$. *Phys. Lett.*, B512:277–282, 2001.
- [7] J. M. Link et al. Search for CP violation in the decays $D^+ \rightarrow K_S^0 \pi^+$ and $D^+ \rightarrow K_S^0 K^+$. *Phys. Rev. Lett.*, 88:041602, 2002.
- [8] J. M. Link et al. A new measurement of the Ξ_c^+ lifetime. *Phys. Lett.*, B523:53–59, 2001.
- [9] J. M. Link et al. Measurement of natural widths of Σ_c^0 and Σ_c^{++} baryons. *Phys. Lett.*, B525:205–210, 2002.
- [10] J. M. Link et al. A high statistics measurement of the Λ_c^+ lifetime. *Phys. Rev. Lett.*, 88:161801, 2002.
- [11] J. M. Link et al. Evidence for new interference phenomena in the decay $D^+ \rightarrow K^- \pi^+ \mu^+ \nu$. *Phys. Lett.*, B535:43–51, 2002.
- [12] J. M. Link et al. New measurement of the D^0 and D^+ lifetimes. *Phys. Lett.*, B537:192–200, 2002.
- [13] J. M. Link et al. Measurements of relative branching ratios of Λ_c^+ decays into states containing Σ . *Phys. Lett.*, B540:25–32, 2002.
- [14] J. M. Link et al. A new measurement of the Ξ_c^0 lifetime. *Phys. Lett.*, B541:211–218, 2002.

- [15] J. M. Link et al. Measurement of the D^+ and D_s^+ decays into $K^+K^-K^+$. *Phys. Lett.*, B541:227–233, 2002.
- [16] J. M. Link et al. New measurements of the $\Gamma(D^+ \rightarrow \bar{K}^{*0}\mu^+\nu)/\Gamma(D^+ \rightarrow K^-\pi^+\pi^+)$ and $\Gamma(D_s^+ \rightarrow \phi\mu^+\nu)/\Gamma(D_s^+ \rightarrow \phi\pi^+)$ branching ratios. *Phys. Lett.*, B541:243–250, 2002.
- [17] J. M. Link et al. New measurements of the $D^+ \rightarrow \bar{K}^{*0}\mu^+\nu$ form factor ratios. *Phys. Lett.*, B544:89–96, 2002.
- [18] J. M. Link et al. Observation of a 1750 MeV/ c^2 enhancement in the diffractive photoproduction of K^+K^- . *Phys. Lett.*, B545:50–56, 2002.
- [19] J. M. Link et al. Study of the Cabibbo-suppressed decay modes $D^0 \rightarrow \pi^-\pi^+$ and $D^0 \rightarrow K^-K^+$. *Phys. Lett.*, B555:167–173, 2003.
- [20] J. M. Link et al. Charm system tests of CPT and Lorentz invariance with FOCUS. *Phys. Lett.*, B556:7–13, 2003.
- [21] J. M. Link et al. Measurement of the Ω_c^0 lifetime. *Phys. Lett.*, B561:41–48, 2003.
- [22] J. M. Link et al. Study of hadronic five-body decays of charmed mesons. *Phys. Lett.*, B561:225–232, 2003.
- [23] J. M. Link et al. Studies of correlations between D and \bar{D} mesons in high energy photoproduction. *Phys. Lett.*, B566:51–60, 2003.
- [24] J. M. Link et al. Measurements of Ξ_c^+ branching ratios. *Phys. Lett.*, B571:139–147, 2003.
- [25] J. M. Link et al. Search for rare and forbidden 3-body di-muon decays of the charmed mesons D^+ and D_s^+ . *Phys. Lett.*, B572:21–31, 2003.
- [26] J. M. Link et al. Study of the decay mode $D^0 \rightarrow K^-K^-K^+\pi^+$. *Phys. Lett.*, B575:190–197, 2003.
- [27] J. M. Link et al. Charm-anticharm baryon production asymmetries in photon-nucleon interactions. *Phys. Lett.*, B581:39–48, 2004.
- [28] J. M. Link et al. Dalitz plot analysis of D_s^+ and D^+ decay to $\pi^+\pi^-\pi^+$ using the K-matrix formalism. *Phys. Lett.*, B585:200–212, 2004.
- [29] J. M. Link et al. Measurement of masses and widths of excited charm mesons D_2^* and evidence for broad states. *Phys. Lett.*, B586:11–20, 2004.

- [30] J. M. Link et al. Measurements of six-body hadronic decays of the D^0 charmed meson. *Phys. Lett.*, B586:21–26, 2004.
- [31] J. M. Link et al. New measurements of the $D_s^+ \rightarrow \phi \mu^+ \nu$ form factor ratios. *Phys. Lett.*, B586:183–190, 2004.
- [32] J. M. Link et al. Study of hadronic five-body decays of charmed mesons involving K_S^0 . *Phys. Lett.*, B586:191–197, 2004.
- [33] J. M. Link et al. Measurement of the ratio of the vector to pseudoscalar charm semileptonic decay rate $\Gamma(D^+ \rightarrow K^{*0} \mu^+ \nu_\mu) / \Gamma(D^+ \rightarrow K^0 \mu^+ \nu_\mu)$. *Phys. Lett.*, B598:33–41, 2004.
- [34] J. M. Link et al. Study of the doubly and singly Cabibbo suppressed decays $D^+ \rightarrow K^+ \pi^+ \pi^-$ and $D_s^+ \rightarrow K^+ \pi^+ \pi^-$. *Phys. Lett.*, B601:10–19, 2004.
- [35] J. M. Link et al. Measurement of the branching ratio of the decay $D^0 \rightarrow \pi^- \mu^+ \nu$ relative to $D^0 \rightarrow K^- \mu^+ \nu$. *Phys. Lett.*, B607:51–58, 2005.
- [36] J. M. Link et al. A study of $D^0 \rightarrow K_S^0 K_S^0 X$ decay channels. *Phys. Lett.*, B607:59–66, 2005.
- [37] J. M. Link et al. Analysis of the semileptonic decay $D^0 \rightarrow \bar{K}^0 \pi^- \mu^+ \nu$. *Phys. Lett.*, B607:67–77, 2005.
- [38] J. M. Link et al. Measurements of the q^2 dependence of the $D^0 \rightarrow K^- \mu^+ \nu$ and $D^0 \rightarrow \pi^- \mu^+ \nu$ form factors. *Phys. Lett.*, B607:233–242, 2005.
- [39] J. M. Link et al. Study of the decay mode $D^0 \rightarrow K^- K^+ \pi^- \pi^+$. *Phys. Lett.*, B610:225–234, 2005.
- [40] J. M. Link et al. Measurement of the doubly Cabibbo suppressed decay $D^0 \rightarrow K^+ \pi^-$ and a search for charm mixing. *Phys. Lett.*, B618:23–33, 2005.
- [41] J. M. Link et al. Hadronic mass spectrum analysis of $D^+ \rightarrow K^- \pi^+ \mu^+ \nu$ decay and measurement of the $K^*(892)^0$ mass and width. *Phys. Lett.*, B621:72–80, 2005.
- [42] J. M. Link et al. A measurement of the D_s^+ lifetime. *Phys. Rev. Lett.*, 95:052003, 2005.

- [43] J. M. Link et al. Search for a strongly decaying neutral charmed pentaquark. *Phys. Lett.*, B622:229–238, 2005.
- [44] J. M. Link et al. Search for T violation in charm meson decays. *Phys. Lett.*, B622:239–248, 2005.
- [45] J. M. Link et al. Study of Λ_c^+ Cabibbo favored decays containing a Λ baryon in the final state. *Phys. Lett.*, B624:22–30, 2005.
- [46] J. M. Link et al. Search for $\Lambda_c^+ \rightarrow p K^+ \pi^-$ and $D_s^+ \rightarrow K^+ K^+ \pi^-$ using genetic programming event selection. *Phys. Lett.*, B624:166–172, 2005.
- [47] J. M. Link et al. A non-parametric approach to the $D^+ \rightarrow \overline{K}^{*0} \mu^+ \nu$ form factors. *Phys. Lett.*, B633:183–189, 2006.
- [48] J. M. Link et al. Study of the decay asymmetry parameter and CP violation parameter in the $\Lambda_c^+ \rightarrow \Lambda \pi^+$ decay. *Phys. Lett.*, B634:165–172, 2006.
- [49] J. M. Link et al. New measurement of $\text{BR}(D^+ \rightarrow \rho^0 \mu^+ \nu) / \text{BR}(D^+ \rightarrow \overline{K}^{*0} \mu^+ \nu)$ branching ratio. 2005. hep-ex/0511022, accepted by *Phys. Lett. B*.
- [50] J. M. Link et al. Reconstruction of vees, kinks, Ξ^- 's, and Ω^- 's in the FOCUS spectrometer. *Nucl. Instrum. Meth.*, A484:174–193, 2002.
- [51] J. M. Link et al. Čerenkov particle identification in FOCUS. *Nucl. Instrum. Meth.*, A484:270–286, 2002.
- [52] J. M. Link et al. The target silicon detector for the FOCUS spectrometer. *Nucl. Instrum. Meth.*, A516:364–376, 2003.
- [53] J. M. Link et al. Application of genetic programming to high energy physics event selection. *Nucl. Instrum. Meth.*, A551:504–527, 2005.